

Neoplasia: The process of abnormal and uncontrolled growth of cells. The product of neoplasia is a **neoplasm** (a tumor).

Science of the Total Environment 409 (2011) 3649–3665



Contents lists available at ScienceDirect

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv



Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil [☆]

Adilza C. Dode ^{a,b,e,*}, Mônica M.D. Leão ^c, Francisco de A.F. Tejo ^d, Antônio C.R. Gomes ^e, Daiana C. Dode ^{e,f}, Michael C. Dode ^e, Cristina W. Moreira ^b, Vânia A. Condessa ^b, Cláudia Albinatti ^b, Waleska T. Caiaffa ^g

^a Minas Methodist University Center Izabela Hendrix, Belo Horizonte City, Minas Gerais State, Brazil

^b Municipal Government of Belo Horizonte, Municipal Health Department, Belo Horizonte City, Minas Gerais State, Brazil

^c UFMG—Universidade Federal de Minas Gerais—Belo Horizonte, Environmental and Sanitary Engineering Department, Belo Horizonte City, Minas Gerais State, Brazil

^d UFGC—Universidade Federal de Campina Grande, Center of Electrical Engineering and Informatics, Academic Unit of Electrical Engineering, Paraíba State, Brazil

^e MRE Engenharia (Electromagnetic Radiations Measurement—Engineering), Belo Horizonte City, Minas Gerais State, Brazil

^f Faculty of Medical Sciences, Medicine—Belo Horizonte, Belo Horizonte City, Minas Gerais State, Brazil

^g UFMG—Universidade Federal de Minas Gerais—Belo Horizonte, Urban Health Observatory, Belo Horizonte City, Minas Gerais State, Brazil

ARTICLE INFO

Article history:

Received 14 January 2011

Received in revised form 25 May 2011

Accepted 25 May 2011

Available online 13 July 2011

Keywords:

Assessment and management of impacts and environmental risks

Non-ionizing electromagnetic radiation

Public health

Radio base station

Environmental electromagnetic pollution

Environmental electromagnetic

field monitoring

ABSTRACT

BS = cellular antenna Base Station

Pollution caused by the electromagnetic fields (EMFs) of radio frequencies (RF) generated by the telecommunication system is one of the greatest environmental problems of the twentieth century. The purpose of this research was to verify the existence of a spatial correlation between base station (BS) clusters and cases of deaths by neoplasia in the Belo Horizonte municipality, Minas Gerais state, Brazil, from 1996 to 2006 and to measure the human exposure levels to EMF where there is a major concentration of cellular telephone transmitter antennas. A descriptive spatial analysis of the BSs and the cases of death by neoplasia identified in the municipality was performed through an ecological–epidemiological approach, using georeferencing. The database employed in the survey was composed of three data banks: 1. death by neoplasia documented by the Health Municipal Department; 2. BSs documented in ANATEL (“Agência Nacional de Telecomunicações”: “Telecommunications National Agency”); and 3. census and demographic city population data obtained from official archives provided by IBGE (“Instituto Brasileiro de Geografia e Estatística”: “Brazilian Institute of Geography and Statistics”). The results show that approximately 856 BSs were installed through December 2006. Most (39.60%) of the BSs were located in the “Centro-Sul” (“Central-Southern”) region of the municipality. Between 1996 and 2006, 7191 deaths by neoplasia occurred and within an area of 500 m from the BS, the mortality rate was 34.76 per 10,000 inhabitants. Outside of this area, a decrease in the number of deaths by neoplasia occurred. The greatest accumulated incidence was 5.83 per 1000 in the Central-Southern region and the lowest incidence was 2.05 per 1000 in the Barreiro region. During the environmental monitoring, the largest accumulated electric field measured was 12.4 V/m and the smallest was 0.4 V/m. The largest density power was 40.78 $\mu\text{W}/\text{cm}^2$, and the smallest was 0.04 $\mu\text{W}/\text{cm}^2$.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Mobile phone radio base stations (RBSs) are now found in cities and communities worldwide. They can be found near or even on top of homes, schools, hospitals, daycare centers and offices. In Brazil, the number of mobile phone users is estimated to be over 200 million and there are more than 5 billion users worldwide. In the municipality of Belo Horizonte, the capital of the state of Minas Gerais, there are approximately 1000 base stations (BSs) with 128.77 accesses by

mobile phones per 100 inhabitants and in Brazil, there are 49,979 BSs licensed through April 2011 (ANATEL, 2011).

The non-ionizing electromagnetic radiation from the BSs is of low intensity compared to the current guidelines on human exposure limits. However, its emission is continuous. This raises concerns as to whether the health and well-being of people living or working close to the BSs are at risk (Khurana et al., 2010; Alanko et al., 2008).

The emission of a BS is usually described by its effectively radiated power in watts (W), which describes the total amount of radiation emitted by the antenna of the BS. Their intensity, called the power density, is commonly measured in milliwatts per square centimeter (mW/cm^2) or microwatt per square centimeter ($\mu\text{W}/\text{cm}^2$) and it expresses the power per unit area impinging normally to the external surface of the subject. The immission (absorption) of the subject is measured by the specific absorption rate (SAR), which is reported in

[☆] All the authors declare that they have no conflicts of interest.

* Corresponding author at: Rua Desembargador Assis Rocha, 279, Bairro Belvedere, 30320-250, Belo Horizonte (MG), Brazil. Tel.: +55 031 3286 1899.

E-mail addresses: adilzadode@terra.com.br (A.C. Dode), monica@desa.ufmg.br (M.M.D. Leão).

watts per kilogram of body tissue (W/kg). The SAR reflects the power that is locally absorbed in a certain volume of biological tissue and is proportional to the square of the local magnitude of the electric field intensity. For ethical reasons, the SAR can only be assessed on animal models or inferred from virtual (computational) models of animal or human subjects (Lai, 2000).

Some scientific studies have shown evidence of increased numbers of cancer cases for people living less than 500 m from the BSs (Eger et al., 2004; Wolf and Wolf, 2004; Eger and Jahn, 2010).

In the Belo Horizonte municipality and in many other urbanized cities and communities in Brazil, the mobile phone network is deployed in regions of high demographic density close to homes and on the facades and roofs of public or private buildings. It is also common to have several antennas sharing the same support structure.

This situation motivated the research of Dode (Dode, 2003) in the Belo Horizonte municipality, where a methodology designed to assess the levels of electromagnetic radiation exposure of the dwellers was used, based on the technical specifications of a sample of the installed BSs. Those estimated data were then compared to measured *in situ* data for the same set of BSs. Fig. 1 illustrates the site of a typical BS (Base Station BH 20) in a residential area of the Serra neighborhood, in the Belo Horizonte municipality and Fig. 2 shows its geographical location. Fig. 3 represents the horizontal and vertical radiation patterns per sector of the same BS. This diagram has been obtained from the technical archives documented by the operators in the Secretaria Municipal de Meio Ambiente (Municipal Environmental Department), the official organization of the municipality that is responsible for the environmental licensing of the BSs.

Some studies have shown evidence of general risks to health and specific risks of cancer associated with the physical proximity of the transmitter antennas of the telecommunication network.

One of the first of these studies indicated an association between cancer growth and a residence near a transmitter antenna (Cherry, 1999). Later, Santini et al. (Santini et al., 2002) carried out a qualitative survey of 530 people living within 300 m of a certain BS. Despite the

subjective methodology, the study showed a peak of symptoms occurred at locations in the interval between 50 and 100 m from the BS, which coincided with the typical distances at which the main lobe reached the ground. In another study, also in France, Santini et al. (2003) surveyed dwellers living 300 m from the BS and others who lived farther away and found more complaints about irritability, depressive tendency, memory loss, problems with concentration, dizziness, within 100 m; headache, sleeping disorder, discomfort, skin problem within 200 m; and fatigue within 300 m, among those living closer to the BS and showed more variability in disease occurrence with distance. Again, this study contained some biases because it was subjective and therefore did not result in a conclusion about the relationship between cancer and the amount of radiation exposure.


Navarro et al. (2003) conducted a study of 145 people in Múrcia, Spain, but only included 101 questionnaires in the analysis. Two groups of participants were formed: one that was living within 150 m of the BS and another beyond 150 m. The average measured power density was $1.1 \mu\text{W}/\text{cm}^2$ at locations within 150 m and $0.1 \mu\text{W}/\text{cm}^2$ beyond 150 m. This study also showed that complaints (insomnia, headaches, difficulty in concentration, discomfort) were greater at locations where the power density was higher, inside the 150 m range.

In Poland, Gadzicka (Gadzicka et al., 2006) also used a questionnaire to conduct a neurobehavioral clinical study involving 500 subjects. The most important finding was the incidence of frequent headaches in subjects living less than 150 m from the BS. However, the study was limited because of the lack of information about the technical characteristics of the BS and the measurements of electromagnetic exposure.

Eger et al. (2004) carried out research in the city of Naila, Germany, to examine whether people who live near mobile phone BSs were at any risk of becoming ill with malignant tumors. Their data bank consisted of records of patients from 1994 to 2004. While preserving the privacy of the information, the personal data of almost 1000 subjects were examined. The analysis showed that the number of newly developed cancer cases was significantly higher among those



Fig. 1. BS site BH 20 in a residential area of the Serra neighborhood in Belo Horizonte municipality.



This is the pattern of a sector antenna. You need 4 for 360 degree spread.

KATHREIN	Date 21 06 1995	Horizontal and Vertical Radiation Patterns 870 MHz	Type 737656
	Name		

Fig. 3. Horizontal and vertical radiation patterns per sector of BS site BH 20 (KATHREIN MOBILCOM BRASIL LTDA, HUEMER E, and LENSIG KI-, 1999).

incidence in the women of group A was significantly higher ($p < 0.0001$) than the cancer incidence of group B and the city as a whole. A relative risk comparison revealed that there were approximately 4.15 more cases of cancer in group A than in the population as a whole. The results, although still not conclusive, indicated a necessity to revise the current exposure limits in favor of more protective levels. Both the estimated and measured power densities in the entire exposed area in Netanya were far below $0.53 \mu\text{W}/\text{cm}^2$, that is, approximately 800 times lower than the exposure limit of $425 \mu\text{W}/\text{cm}^2$ for the frequency of 850 MHz from the ICNIRP guidelines.

The aforementioned studies, which aimed to find evidences of an increase in cancer incidence with proximity to mobile phone BSs, warrant additional research, because the cellular phone technology is relatively new and the associated total amount of environmental radiation is far from negligible.

The inhabitants of the Belo Horizonte municipality and the scientific community in general are also concerned about the number

of already installed BSs and the proliferation of new wireless BSs, not only for telephony but also for television. The number of mobile phone BSs, which equaled 474 in 2003, had reached approximately 856 in 2006.

Thus, this research to study health was conducted in a broad environmental context, aiming to verify if there is a spatial correlation between the cellular telephony system BS location and the cases of death by neoplasia during the period between 1996 and 2006.

2. Materials and methods

2.1. Area of study

The Belo Horizonte municipality, with an area of approximately 300 km^2 of area, has a tropical climate and is located at an average altitude of 900 m (minimum of 800 m and maximum of 1200 m)




Fig. 4. The Belo Horizonte municipality and the nine SDs.

above sea level. The geology of the city includes several kinds of minerals and its soil of crystalline rocks is composed of dolomite, quartzite, phyllite and various schists. Constructed over many hills, the city is surrounded by a mountain named "Serra do Curral". The municipality is divided into nine regions or sanitary and administrative districts (SDs): "Centro-Sul" ("Central-Southern"), "Norte" ("Northern"), "Leste" ("Eastern"), "Oeste" ("Western"), "Noroeste" ("Northwestern"), "Nordeste" ("Northeastern"), "Venda Nova", "Pampulha" and "Barreiro". Fig. 4 shows the Belo Horizonte municipality and the nine SDs.

The municipality has 55 universities and colleges, 36 hospitals, and a subway system containing 19 stations and 29 km of track, which transports 145,000 passengers a day. The majority of households are served by potable water (99.3%), garbage collection (96.6%), sewage (93.2%), electric power (99.83%), and landline phones (81.43%) (UNDP, 2008) and 128.77 accesses by mobile phones per 100 inhabitants through April 2011 (ANATEL, 2011). With these data, one can conclude that much of the population possesses a cellular phone and more than 28% of the inhabitants have more than one. The city has road and railroad networks that link it to the main centers of the country, as well as three airports.

More than 80% of the municipal economy is focused on commerce, financial services, real estate activities and public administration. The metallurgical industries, including iron and steel metallurgy, and ore mining are located in the areas surrounding the metropolitan region but not in the Central-Southern region.

The city of Belo Horizonte has been selected by the Population Crisis Committee of the United Nations (UN, 2007) as the metropolis with the best quality of life in Latin America and was ranked 45th in the world. Its health system is considered very good, according to the Atlas of Human Development (2000)/United Nations Development Programme (UNDP, 2008).

The city had 2,238,332 inhabitants in 2003 and 2,258,096 in 2010 (IBGE, 2010), which suggests that the population in the city is stable. However, the city, as in any urban area in Brazil, has a concentrated

population, with a large number of people living in apartment buildings. This fact, along with the mountainous landscape, force mobile phone operators to install their BSs at strategic points in the city, mainly on top of towers and poles, as on the terraces of public or residential buildings, to ensure good coverage of the mobile phone network.

Of the nine SDs, the Central-Southern region is the richest region of the city and is the third largest in number of inhabitants with 249,862. There are plenty of commercial and service shops, several shopping centers, and many households with one or more families. This SD also has several hospitals, parks and leisure areas. Most of the dwellers are highly educated and belong to the middle and upper classes. The traffic is heavy because of the large number of vehicles that travel in that region. The Western region is less densely populated, has no skyscrapers and its inhabitants have low revenues. The Barreiro region is the most populated after the Central-Southern SD, and has many industries. The most populated region, with 338,753 inhabitants, is the Northwestern (IBGE, 2000).

2.2. Study design

This ecological study consists of an exploratory spatiotemporal analysis to determine whether there is an association between clusters of BSs and deaths by neoplasia in the Belo Horizonte municipality, in the southeastern part of Brazil. This design was chosen because of the possibility of using geographic areas as units of analysis, where each unit of analysis is composed of a group of individuals or communities. Therefore, it is possible to determine whether there is a correlation between a certain risk and the occurrence of certain grievances within the population. In this type of study, it is not possible to consider individual characteristics, such as food and life habits, activity level, smoking, self-medication, individual pathologies, or genetic factors (GORDIS, 2004).

The analysis was based on the following databases: 1. A database of deaths by neoplasia documented in the Mortality Information

System (SIM: "Sistema de Informação em Mortalidade"), provided by the City Health Department; 2. A database of the site register of BSs, provided by the Brazilian Telecommunications Agency (ANATEL); 3. A database of the city census, including demographic information provided by the Brazilian Institute of Geography and Statistics (IBGE).

The death, BS and population data were geocoded according to census tracts (CTs) or censitarian sectors (CSs), which are "territorial units defined by IBGE (IBGE, 2000) to orient the spatial distribution of a population". The definition of a CT is related to a specific geographical zone whose population can be counted by local

interviewers, taking into account the existence of geographical barriers, the population size and traffic flow. There were a total of 2563 CTs in the Belo Horizonte municipality (IBGE, 2000).

2.2.1. Cancer death variable

The main outcome that was studied was the number of deaths by neoplasia of Belo Horizonte municipality residents that occurred from 1996 to 2006, were reported to the City Health Department and were routinely confirmed by established criteria, under the responsibility of

Table 1
International classification of diseases – ICD-10.

Disease	ICD-10 – According to WHO ICD10 homepage		Bibliographical references
	Primary	Secondary	
Primary: Malignant melanoma of skin; Other malignant neoplasms of skin./Secondary malignant neoplasm of skin.	C43 and C44	C79.2	Eger et al., 2004.
Primary: Malignant neoplasm of breast./Secondary malignant neoplasm of other specified sites.	C50	C79.8	Eger et al., 2004; Wolf and Wolf, 2004; Bioinitiative Report, 2007; Guenel et al., 1996; Feychting et al., 1997; Wakeford, 2004; Mack et al., 1991; Beall et al., 1996; Beniashvili et al., 2005; Hardell and Sage, 2007.
Primary and secondary: Malignant neoplasm without specification of site.	C80	C80	Khurana, 2008; Hardell et al., 2007; Bioinitiative Report, 2007; Mack et al., 1991; Beall et al., 1996; Guenel et al., 1996; Wakeford, 2004.
Primary: Malignant neoplasm of ovary./Secondary malignant neoplasm of ovary.	C56	C79.6	Eger et al., 2004; Wolf and Wolf, 2004
Primary: Hodgkin's Disease./Secondary and unspecified malignant neoplasm of lymph nodes.	C81	C77	Wolf and Wolf, 2004.
Primary: Malignant neoplasm of bronchus and lung./Secondary malignant neoplasm of lung	C34	C78.0	Eger et al., 2004; Wolf and Wolf, 2004
Primary: Malignant neoplasm of kidney, except renal pelvis./Secondary malignant neoplasm of other sites.	C64	C79.0	Wolf and Wolf, 2004.
Primary: Malignant neoplasm of prostate./Secondary malignant neoplasm of other specified sites.	C61	C79.8	Eger et al., 2004.
Primary: Malignant neoplasm of pancreas; Pancreas, unspecified./Secondary malignant neoplasm of other and unspecified digestive organs.	C25 and C25.9	C78.8	Eger et al., 2004.
Primary: Malignant neoplasm of other and ill-defined digestive organs: Intestinal tract, part unspecified; Malignant neoplasm of small intestine; Malignant neoplasm of colon; Malignant neoplasm of rectosigmoid junction./Secondary malignant neoplasm of small intestine; Secondary malignant neoplasm of large intestine and rectum.	C26.0; C17; C18; C19	C78.4; C78.5	Eger et al., 2004.
Primary: Malignant melanoma of skin; Melanoma <i>in situ</i> ./Secondary malignant neoplasm of skin	C43 and D03	C79.2	Eger et al., 2004; Hallberg, 2004; Johansson, 2006.
Primary: Malignant melanoma of skin./Secondary malignant neoplasm of skin.	C43	C79.2	Stang, 2001.
Primary: Malignant neoplasm of kidney, except renal pelvis; Malignant neoplasm of renal pelvis./Secondary malignant neoplasm of kidney and renal pelvis.	C64 and C65	C79.0	Eger et al., 2004.
Primary: Malignant neoplasm of stomach./Secondary malignant neoplasm of other and unspecified digestive organs.	C16	C78.8	Eger et al., 2004.
Primary: Malignant neoplasm of bladder./Secondary malignant neoplasm of bladder and other and unspecified urinary organs.	C67	C79.1	Eger et al., 2004.
Primary: Multiple myeloma and malignant plasma cell neoplasms; Lymphoid leukemia; Myeloid leukaemia; Monocytic leukemia; Other leukemias of specified cell type; Leukemia of unspecified cell type; Other and unspecified malignant neoplasms of lymphoid, haematopoietic and related tissue.	C90; C91; C92; C93; C94; C95 and C96		Eger et al., 2004.
Primary: Hodgkin's disease; Follicular [nodular] non-Hodgkin's lymphoma; Diffuse non-Hodgkin's lymphoma; Peripheral and cutaneous T-cell lymphomas; Other and unspecified types of non-Hodgkin's lymphoma./Secondary and unspecified malignant neoplasm of lymph nodes.	C81; C82; C83; C84 and C85	C77	Hardell et al., 2007.
Primary: Malignant neoplasm of brain./Secondary malignant neoplasm of brain and cerebral meninges.	C71	C79.3	Khurana, 2008; Hardell et al., 2007; Schoemaker et al., 2005.

the epidemiology officers of the city, accredited by federal and local health authorities (BRAZIL, 2011).

All deaths by neoplasia, based on death certificates, were provided. Then, they were re-selected according to a subset of the International Classification of Diseases (ICD) previously organized and extracted from a careful review of the scientific literature, linking cancer and non-ionizing electromagnetic radiation, as can be seen in Table 1.

Out of 22,493 deaths that occurred in the analyzed period (1996 to 2006), 7191 were initially eligible for the study. The selected death by neoplasia cases were grouped according to the CT of the residences, based on the residents' postal address. The data bank of SIM did not possess the address of the persons who died by neoplasia in 1998. So, about 780 deaths that occurred in that year could not be georeferenced. To identify the CT, the cartographic map was used, within the borders delimited by IBGE (IBGE, 2000). Fig. 5 shows the fluxogram of deaths by neoplasia in the period from 1996 to 2006.

The death cases were further analyzed according to age, gender, site of residence and year of occurrence and the death rates were determined as described below. After aggregation of the deaths and BS exposure (explained below) in the CT, differing numbers of deaths were determined, depending on whether the date of first exposure was taken to be the date of the first license of the BS (7191 deaths) or the date of the register of the BS (8082 deaths). We opted to work with the date of first license; an option that makes our analysis even more conservative.

2.2.2. Base stations

The BS database and their respective geographical locations were obtained from the ANATEL database (site: <http://www.anatel.gov.br>) and were further geoprocessed according to their CT in two distinct years: 2003 and 2006. In 2003, there were approximately 474 BSs, and in 2006, there were approximately 856 in the city. Clusters (the so-called "hotspots") were the identified in each SD. This explanatory analysis was carried out through thematic maps, using the software MAPINFO™, version 7.0, and the Kernel estimator.

2.2.3. Data processing and mapping of the BSs and deaths in the Belo Horizonte municipality

Eligible deaths by neoplasia were then plotted inside circles with radii varying from 100 m to 1000 m, centered at the location of the first transmitter antenna of the mobile phone network to which the resident was possibly exposed. This selection took into account the date of the death and either the date of registration or of the first

license of the given BS. To detect case conglomerates in space, the total amount (2563) of CTs and the corresponding nine DSs of the city were used again. Software was developed to calculate the shortest distance between the death and the antenna and to estimate the time of exposure of the deaths to the radiation of the antennas.

2.2.4. Death rates

The mortality by neoplasia rates per CT were determined from the neoplasia diagnoses in compliance with the ICD-10 during the period of the study, using the CT as defined by IBGE for the 2000 national census as the spatial analysis unit. The deaths in every CT were used as the numerator. The temporal unit was the calendar year. The death data were then georeferenced to the address of the subject's and the IBGE census data (IBGE, 2000) for each CT. The estimated population at risk was used as the denominator.

The accumulated incidence in the Belo Horizonte municipality was calculated by dividing the total amount of deaths in each region by region's entire population.

2.2.5. Estimates of the mortality rates according to distance and duration of exposure to the radiation of the BSs

To understand the spatiotemporal exposure to the radiation of the BSs, the duration of possible exposure corresponding to each death was estimated, using a proxy to the subject's residential addresses, in terms of the duration of his or her exposure to the first installed transmitter antenna of the mobile phone network. To estimate the number of days of exposure, the elapsed time period from the date of installation of the first antenna to the date of the death was calculated, in spite of the exposure to the radiation of other antennas that might have been installed afterward. Some subjects may have been exposed to many antennas at different times, but in this study we considered only the date of installation of the first antenna. The delimitation of the distance intervals of the BSs was then performed. Each interval with a radius of 100 m had a BS as its center, from which the distance was increased to 1000 m. For each 100-meter interval, the deaths that occurred within the elapsed period of time, as well as the estimated population living within that radius, was then observed. To obtain the population at risk, the estimates of all of the CTs were considered, even those which were only partially included within those radii. Therefore, the population at risk was conservatively overestimated.

To estimate the mortality rate within each radius, the number of deaths was divided by the estimated population included in the radius of each CT. For example, for the 100-meter radius, the 3569 deaths were divided by the 821,890 estimated exposed subjects living inside that radius. For the rates between 200 m and 1000 m, both the number of deaths and the included population were cumulatively considered. This was necessary because the subjects included in the 100-meter radius must be considered to perform the calculation inside the 200-meter radius (Fig. 6) (Table 5).

2.3. Environmental monitoring of the electromagnetic field (EMF)

In 2008, one began monitoring the environmental EMF in the CT with the largest concentration of antennas in the Belo Horizonte municipality. The survey used an electric field meter and isotropic probe, with a frequency range of 0.2 MHz to 3.0 GHz; a spectrum analyzer, with a frequency range of 10.0 MHz to 6.0 GHz; a datalogging multimeter; a GPS unit and a laptop. For the field measurements, the following guidelines were observed: IEEE, 1999, 1992; NCRP, 1993; ANATEL Annex to Resolution no. 303 (ANATEL, 2000a, 2000b), and the environmental field survey of two particular nearby BSs, carried out by Dode (Dode, 2003). The analyzed frequencies of the BSs were corresponded to the bands A, B, C and D. During the measurements, stronger electric field intensities were usually found when the probe was far from the ground. Approximately 400 points in the Central-Southern





Fig. 5. Fluxogram of deaths by neoplasia in the period from 1996 to 2006.



Within 100 meters = 3,569 deaths
Within 200 meters = 3,569 + 1,408 deaths = 4,977 deaths
Within 300 meters = 4,977 + 973 deaths = 5,950 deaths
Within 400 meters = 5,950 + 482 deaths = 6,432 deaths
Within 500 meters = 6,432 + 292 deaths = 6,724 deaths and so on within 1000 meters
Beyond 1000 meters + 147 deaths
 Total amount of = 7,191 deaths

Fig. 6. Total amount of deaths by neoplasia per 100-meter distance band, in census tracts inside a radius of up to 1000 m from the mobile phone transmitter antennas, in the Belo Horizonte municipality, from 1996 to 2006. Total: 7044 deaths.

region of the municipality, which were located in squares, parks, schools and households nearby BSs, were considered in the survey.

2.4. Ethical committees

Because this study includes data on human beings, it was approved by the ethical committees of the Institutional Review Boards of the Federal University of Minas Gerais and the Belo Horizonte City Health Department, with the purpose of accomplishing the Resolution 196/1996 of the Brazilian Health Ministry.

3. Results

3.1. Total deaths by neoplasia selected in the period from 1996 to 2006

Fig. 7 shows the geographic location of the cases of deaths by neoplasia that were confirmed in the literature and selected according to Table 1, totaling 7191 deaths. The detailed geographic description of

deaths location can be seen in Table 2. The Central-Southern SD contained the greatest absolute number of deaths, followed by the Northwestern and Eastern SDs.

3.2. Base station

All registered BSs that were georeferenced through 2006 are plotted in Fig. 8. The percentage of BSs installed through December 2003 was the greatest in the Central-Southern region, comprising 38.60% (182 out of 474); until December 2006, the percentage was approximately 39.60% (338 out of 856). The BS percentage by region in the Belo Horizonte municipality in 2003 and 2006 can be seen in Fig. 9.

3.3. Data processing and mapping of the base stations and deaths in the Belo Horizonte municipality

Fig. 10 portrays a sample of the georeferencing of the BSs and the deaths by neoplasia in downtown Belo Horizonte City located

Table 2
Description of the death coding and geographic location.

Deaths codification	Regions or sanitary districts									Total
	Barreiro	Central-Southern	Eastern	Northeastern	Northwestern	Northern	Western	Pampulha	Venda Nova	
C16	83	143	132	124	183	75	125	57	89	1011
C17	2	3	4	1	3	2	4	4	0	23
C18	1	2	2	2	1	1	3	0	1	13
C19	4	8	13	9	16	2	7	1	5	65
C25	30	155	88	74	137	23	77	42	39	665
C26	4	27	17	12	17	19	23	14	14	147
C34	88	300	194	140	233	89	187	75	99	1405
C43	0	0	0	0	0	0	0	0	1	1
C50	43	210	145	68	177	23	94	35	34	829
C61	42	174	131	84	186	51	122	56	58	904
C64	17	40	28	19	28	16	20	10	11	189
C65	0	1	0	1	0	0	1	0	3	6
C67	18	51	42	28	40	10	30	21	12	252
C71	37	105	54	37	94	23	63	30	28	471
C80	36	86	71	55	83	36	61	21	34	483
C81	5	19	8	4	9	3	9	3	5	65
C83	0	2	0	0	1	0	0	0	0	3
C84	0	0	0	0	0	0	1	0	0	1
C90	11	40	39	21	41	14	33	13	16	228
C91	4	11	6	10	12	3	15	7	7	75
C92	19	62	34	24	43	13	34	14	16	259
C93	0	1	0	0	1	0	0	0	1	3
C94	0	2	0	0	0	0	0	0	0	2
C95	7	17	10	4	18	7	12	4	12	91
Total	451	1459	1018	717	1323	410	921	407	485	7191

in Central-Southern region. A given BS can have three, six, nine, twelve or more antennas, depending on the requirements in the region.

To detect clusters of cases in space, the nine SDs in the Belo Horizonte and their 2563 CTs were used as units of analysis. In Fig. 11, there are CTs with 12, 13, 14 and even 18 deaths.

Table 3
Percentage of deaths by age and gender in Belo Horizonte municipality.

Age	Male	Female	Deaths total	Percentage%
00–04	10	16	26	0.36
05–09	13	10	23	0.32
10–14	12	8	20	0.28
15–19	11	8	19	0.26
20–29	34	34	68	0.95
30–39	80	120	200	2.78
40–49	247	322	569	7.91
50–59	535	559	1,094	15.21
60–69	920	686	1,606	22.33
70–79	1,217	797	2,014	28.00
80–89	708	550	1,258	17.49
90–99	136	158	294	4.08
TOTAL	3,923	3,268	7,191	

Table 4
Accumulated incidence rate of all deaths in the Belo Horizonte municipality.

Regions or sanitary districts	Population	Death number	Accumulated incidence rate/1000
Centro-Sul	249,862	1459	5.83
Leste	251,118	1018	4.05
Noroeste	338,753	1323	3.90
Pampulha	106,330	407	3.82
Oeste	249,059	921	3.69
Nordeste	248,406	717	2.88
Norte	153,821	410	2.66
Venda Nova	198,475	485	2.44
Barreiro	219,873	451	2.05
TOTAL	2,015,697	7191	

3.4. Death rate

The percentage of deaths by neoplasia per year in the Belo Horizonte municipality from 1996 to 2006, considering the start of exposure to be the date of the first license, is shown in Fig. 12. The accumulated incidence rate per 1000 residents for each SD is shown in Table 4. Again the Central-Southern SD presented the highest accumulated incidence rate, i.e., 5.83 incidents per 1000 inhabitants and the lowest rate was 2.05 incidents per 1000 inhabitants in the Barreiro SD.

The same trend was observed for both women and men with similar profiles during the studied years. As expected the incidence of death of women and men was higher in those older than 40, and in this group, the number of deaths was 3923 for men and 3268 for women. After the age of 40, the death rate (7.91%) grew for both sexes, as shown in Table 3. After the age of 60, the rate was even higher (22.33%).

Supplementary Graphics 1 shows the rate of death by neoplasia, according to ICD classification. The most significant causes were malignant neoplasm of bronchus and lung (C34), 19.55%; malignant neoplasm of stomach (C16), 14.05%; malignant neoplasm of prostate

Table 5
Mortality rates by neoplasia in the Belo Horizonte municipality, according to distance from the BS.

Distance (meters)	Deaths total	Population total	Mortality rate/10,000	Relative risk
Until 100	3569	821,890	43.42	1.35
Until 200	4977	1,237,368	40.22	1.25
Until 300	5950	1,602,869	37.12	1.15
Until 400	6432	1,796,604	35.80	1.11
Until 500	6724	1,934,032	34.76	1.08
Until 600	6869	2,030,093	33.83	1.05
Until 700	6947	2,055,325	33.80	1.05
Until 800	6989	2,086,712	33.49	1.04
Until 900	7000	2,107,277	33.21	1.03
Until 1000	7044	2,148,327	32.78	1.00
Null hypothesis	7,191	2,238,332	32.12	1.00

